

A GUIDE TO THE MEDICAL RISK ASSESSMENT FOR EATING DISORDERS

By Professor Janet Treasure (2009)

www.eatingresearch.com

from the Section of Eating Disorders at
the Institute of Psychiatry and the
Eating Disorders Unit at SLaM

Designed for the use with patients with eating disorders:
Outpatients in primary and second care, medical inpatients, general psychiatric inpatients
and eating disorder patients.

People with eating disorders, in particular those with Anorexia Nervosa, are at high risk in terms of their own health and safety. They have the highest mortality of any psychiatric illness. Both their physical state and suicidal behaviours contribute to this risk. Risk to others is less of a concern.

The factors involved in the assessment of risk in people with eating disorders include:

- Medical risk
- Psychological risk
- Psychosocial risk
- Insight/capacity and motivation

A proxy measure for insight/motivation is the response for treatment. If medical risk is high and there is no response to outpatient treatment, it is necessary to measure capacity and consider the use to mental health law.

This Guide aims to help in the understanding of:

1. **The medical risk** - how to assess it, evaluate it and where to refer
2. **The use of the Mental Health Act** in treatment.

MEDICAL RISK

The medical risk arises from a combination of the restrictive behaviours (food and in some cases fluid) and the compensatory behaviours.

Features in history that indicate medical risk are:

- Excess exercise with low weight
- Blood in vomit
- Inadequate fluid intake in combination with poor eating
- Rapid weight loss
- Factors which disrupt ritualised eating habits (journey/holiday/exam)

Body Mass Index (weight/height²) is a proxy measure of medical risk in Anorexia Nervosa (see Maudsley Body-Mass Index Table). Metabolic changes are most problematic if weight control measures such as vomiting and laxative abuse are used. Neither BMI nor blood tests alone are adequate markers of risk. Screening for risk with an examination of muscle strength, blood pressure, pulse rate, peripheral circulation and core temperature is essential.

Limitations of BMI as a risk marker for Anorexia Nervosa:

- Potential for deceit
- Less reliable if rapid change in weight
- Less reliable at extremes of height
- Higher risk for each BMI range for men (taller)
- Children have BMI range which changes developmentally*
- Less reliable if bulimic features
- Less reliable if fluid restriction

- Less reliable if physical comorbidity
- BMI not critical with regards to risks associated with fluid and electrolyte balance.

*In children and adolescents, the cut off for BMI to make the diagnosis is weight and height below the second centile of BMI. It is possible to get centile charts off the web for the USA (the third centile is depicted) - www.cdc.gov/growthcharts. Also Cole and colleagues have produced charts for population norms.

Brief Essential Medical Examination:

We recommend the following for a rapid risk assessment, repeated frequently as necessary:

- BMI
- Blood pressure and pulse rate, lying and standing
- Muscle strength
- Examination of the skin and temperature for those at high risk for dryness
- A full physical examination looking for e.g. infection (note can be with normal temperature) and signs of nutritional deficiency.

Tests for Muscle Strength (see table below for scoring)

1. The stand up/squat test
The patient is asked to squat down on her haunches and is asked to stand up without using her arms as levers if at all possible.
2. The sit up test:
The patient lies flat on a firm surface, such as the floor, and has to sit up without, if possible, using her hands.

Tests for Hydration

The sign to notice is dizziness or faintness standing up from sitting.

Postural drop, i.e. the difference between lying and standing, blood pressure and heart rate.

Investigations

1. Frequent investigations of full blood count and chemistry (FBC, ESR, UE, Cr, CK, Gluc, LFTs) are necessary if:
 - Patients are in a high risk category from a previous assessment
 - They have a BMI <15 or
 - The BMI is less reliable due to features outlined above, or
 - There is a history of purging.
2. ECG is recommended if BMI <15kg/m² and if drugs which have an effect on QT interval are prescribed.
3. Any other appropriate physical investigations pertinent to physical health.

The table on the next page gives values of concern for each part of the assessment and is followed by a management protocol based on risk.

SYSTEM	Test of Investigation	Concern	Alert
Nutrition	BMI	<14	<12
	Weight loss/week	>0.5kg	>1.0kg
	Skin breakdown	<1.0cm	>0.2cm
	Purpuric rash		++
Circulation	Systolic BP	<90	<80
	Diastolic BP	<70	<60
	Postural drop (sit-stand)	>10	>20
	Pulse rate	<50	<40
Musculo-skeletal (squat and sit-up tests)	Unable to get up without using arms for balance	++	
	Unable to get up without using arms as leverage		++
	Unable to sit up without using arms as leverage	++	
	Unable to sit up at all		++
Temperature		<35C	<34.5C
		98.0F	<97.0F
Bone Marrow	WCC	<4.0	<2.0
	Neutrophil count	<1.5	<1.0
	Hb	<11	<9.0
	Acute Hb drop (MCV and MCH raised - no acute risk)		++
	Platelets	<130	<110
Salt/water balance	K+	<3.5	<3.0
	2. Na+	<135	<130
	3. Mg++	0.5-0.7	<.05
	4. PO4--	0.5-0.8	<0.5
	5. Urea	>7	<10
Liver	Bilirubin	>20	>40
	Alkpase	>110	>200
	AsT	>40	>80
	ALT	>45	>90
	GGT	>45	>90
Nutrition	Albumin	<35	<32
	Creatinine Kinase	>170	>250
	Glucose	<3.5	<2.5
Differential Diagnosis	TFT, ESR		
ECG	Pulse rate	<50	<40
	Corrected QT interval (QTC)		>450msec
	Arrhythmias		++

- The baselines for these tests vary between labs. Any abnormal results is an indication for concern and monitoring.
- A tachycardia in the presence of signs and investigations of severe risk may be a harbinger of imminent cardiovascular collapse.

1. Scores that do not fall into the risk areas

Stable. Regular review and monitoring of the above parameters with routine referral to eating disorders unit/secondary services depending on local resource.

Unstable. If weight is falling, ask the person with Anorexia Nervosa to come up with plan to ensure that nutritional state does not fall into the risk areas. Regularly review the implementation of this plan.

2. Score/s in the concern area

Regular review of parameters (c. weekly) and assessment of capacity with urgent referral to specialised eating disorders team and appropriate medical intervention if needed. As this signifies medical risk, this should also be shared with the carer.

3. Score/s in the alert area

Immediate contact and referral to eating disorders unit and physicians if outpatient with assessment of capacity. The patient will need urgent speciality and medical assessment. If inpatient - immediate contact with on-call physicians.

Useful Tips

Potassium

This is often chronically low in purging, even down to values $<1.5\text{mmol/L}$, with no immediate sequelae. Acute changes are more dangerous. Regular feeding with control of purging is usually sufficient for re-establishment of normal levels. If potassium replacement is required, because it is usually caused by a loss of gastric secretion, it should be done with oral replacement with a salt and water replacement, such as diorlyte, with regular electrolyte review and examination of fluid and water status (measurement of urea and lying and standing blood pressure).

Refractory hypokalaemia can also be due to concurrent low magnesium or calcium and these levels may need checking and rectifying. Serum potassium levels may remain low even with potassium supplements if vomiting persists. A proton pump inhibitor, such as Lansoprazole to inhibit gastric acid secretion, may reduce metabolic alkalosis and help to conserve potassium, but should be a second line measure.

Phosphate

Rebound hypophosphataemia can occur in initial refeeding as it is sequestered by carbohydrate metabolism. It can be lethal. Initial refeeding, including foods with high phosphorus content - e.g. milk-based products ($>2\text{pints/day}$) may be helpful. If necessary, about 4 days of oral phosphate supplementation may also be needed.**

**in Anorexia Nervosa, it is rare for there to be an isolated deficiency of any mineral or vitamin and, therefore, multivitamin and mineral replacement is to be recommended, e.g. Forceval 2 capsules a day or Sanatogen Gold.

Refeeding Oedema

Peripheral oedema is common and harmless during initial refeeding. It resolves within a few weeks spontaneously and rarely needs treatment. It must, however, be distinguished from oedema secondary to heart failure.

Dehydration

This can rapidly lead to medical crisis through circulatory and renal failure. All patients should be fully assessed for dehydration. Take a corroborative history of fluid intake and signs of decompensation (dizziness/ fainting). The physical examination should include assessment of skin turgidity, ocular pressure and lying and standing blood pressure. Regular electrolytes should be checked for high urea, creatinine, sodium and potassium levels. Oral replacement is preferable.

Bradycardia

Investigation: ECG: look for heart block or prolonged QT. Measure U&E. if <40 , admit. Rewarm (if hypothermic) and give a can of Ensure. Monitor HR overnight.

THE USE OF THE MENTAL HEALTH ACT (MHS) IN TREATMENT OF ANOREXIA NERVOSA

Mental Health Act, Mental Capacity Act and Medical Intervention:

If a patient is at medical risk, but does not consent to treatment, they must be:

1. Assessed for capacity
2. Treated under the appropriate legal criteria

Capacity is related to a patient's ability to:

- Understand information relevant to the specific decision;
- Be able to understand the nature of their illness and understand the implications of non-treatment;
- Be able to rationally weigh up the pros and cons of treatment;
- Thus make an informed decision regarding their management and communicate their decision

Under the New MCA (2005), people lacking capacity may be treated if it is in their best interest, if it is the least restrictive option, if it is not depriving them of their liberty and there is no advanced refusal or objection by a donee or court of protection.

Treatment of people with severe Anorexia Nervosa who are not consenting to treatment for their mental disorder will, in most cases, require use of the Mental Health Act as it involves deprivation of liberty and compulsory refeeding. (See MCA 2005 and Deprivation of Liberty Safeguards). Treatment under the Mental Health Act can be given under Section 2 or 3, if they meet the legal criteria (see Code of Practice) whether or not they lack capacity. If after three months from the start of their detention they either lack capacity or object to treatment, SOAD authorisation is required.

For people lacking capacity, emergency medical treatment can be performed under Common Law. Non-emergency treatment for a physical condition not related to the eating disorder may be performed under the MCA 2005.

The laws regarding treatment for 16-18 year olds have changed and for somebody refusing treatment, parental consent cannot be used as authority to treat (see Mental Health Act Code of Practice 2007).

References:

Body Mass Index cut offs to define thinness in children and adolescents: international survey.
Tim J Cole, Katherine M Flegal, Dasha Nicholls and Alan A Jackson
BMJ 2007; 335 (7612): 194